



Ethnobotanical study of wild flora of Haroonabad, District Bahawalnagar, Punjab, Pakistan.

*Zobia Anwer
Siddra Shabbir
Qurat-ul-ain
Tanzeela Iram
Sumaira Tariq
Hina Murad*



Ethnobotanical study of wild flora of Haroonabad, District Bahawalnagar, Punjab, Pakistan

Zobia Anwer¹, Siddra Shabbir², Qurat-ul-ain², Tanzeela Iram², Sumaira Tariq², Hina Murad²

¹Department of Botany, The Islamia University Bahawalpur, Pakistan

²Department of Botany, University of Agriculture Faisalabad, Pakistan

Corresponding Author's Email: toqeerfqw@gmail.com

ABSTRACT

Purpose: The first purpose of this study to record the new and rare use of medicinal plants in the selected area; secondly to discover the plants which were unexplored in the past and thirdly to record the Ethnobotanical data of occurring plant species.

Methodology: The studied area surveyed from July to December. The plants were dried and mounted on standard herbarium sheets. The Ethnobotanical data were collected from 85 local people of Haroonabad by questionnaire.

Findings: The wild flora contained 81 species within 28 families. The largest family was Poaceae with 15 species followed by Euphorbiaceae with 8 species while Asteraceae and Amaranthaceae with 7 species. The life span of plants was comprised of 47 annual species (58%) and 34 perennial species (42%). The life-form spectrum explains that Therophyte 48 species (59%) were the dominant followed by Phanerophytes 7 species (8%) and Chaemophytes 13 species (17%), Hemicryptophyte 8 species (10%), Geophytes 3 species (4%) Halophyte 1 specie (1%) and Parasite 1 specie (1%). Leaf venation classes of plants consisted of reticulate 28 species (34%), pinnate 29 species (36%), parallel 20 species (25%), palmate 2 species (2.5%) and 2 species (2.5%) were leafless thus had no leaf venation. The plant species with herbaceous stems was 68 while with woody stems 13 species. The Ethnobotanical data of 70 species were recorded because 11 plants were not known by anyone. The range of UV value was 0.09 (*Chrozophora plicata* Vahl.) to 0.78 (*Acacia nilotica* L.) and the RCF value range was 0.03 (*Imperata cylindrica* L.) to 0.95 (*Acacia nilotica* L.). The range of ICF value was recorded from 0.45 (Hormonal disorder) to 0.78 (Respiratory disease). 57% whole plant followed by leaves (53%) was used in the ayurvedic field. Therophytes were supported in the studied region for the reason that the region is a semi-arid zone of Punjab, Pakistan.

Unique contribution: The present comprehensive study provides a basic point for other researchers and enhances the knowledge of poor people related to ayurvedic field. The Ethnobotanical study tries to attract the attention towards the conservation strategies of wild plants.

Key words: Survey, Wild flora, Questionnaire, Side effects, Ethnobotanical study

Abbreviations: UV, Use Value; RCF, Relative Frequency Citation; ICF, Informant Consensus Factor

INTRODUCTION

The Ethnobotanical study plays an important role in exposing the relationship between humans and plants [16]. The history of getting medication from the plant is very old. After fulfilling the basic needs like food, fuel shelter and man also uses plants to treat various ailments [13]. In the era we are living in today, 80% of the population is not able to get more expensive treatments. People get medicines from plants to cure their illnesses because herbal medicines are easily available and cheaper than allopathic medicines [1]. There is no doubt that the poor people around the world treat their diseases with herbal medicines. Some plants are specific to the treatment of only one disease and some plants are used to treat many ailments [17]. Ethnobotanical knowledge is now at risk because this knowledge is not transferring from the older generation to the younger generation [2]. The record of the flora of any area by plant taxonomist gain more importance around the world. It provides information about natural vegetation of specific areas [9]. Wild flora in the medicinal field is too much important because, it used in medicinal production and produces immunity in the body to fight against many diseases like cancer, diabetes and heart disease etc. [15]. The purpose of this research to inform the people of developing countries and native people of Haroonabad that wild plants are very important. These wild plants can be used to treat many ailments that are much cheaper than allopathic medicines and their side effects will be much less.

MATERIALS AND METHODS

The whole study area Haroonabad surveyed from July to December. The plants were collected after conducting 15 surveys of the selected area. The plant specimens collected from the area. The specimens were dried and mounted on standard herbarium sheets. The collected specimens were identified with the help of various monographs {(<http://www.ipni.org>), (www.theplantlist.org) and (<http://www.ars-grin.gov/cgi-bin/npgs/html/queries.pl>)}

Climate of studied area: The climate in this area is very hot and harsh. Temperature fluctuations are also very high. In the summer, the mean temperature is between 35°C and 60°C and in winter the mean temperature is between 10°C and 20°C. In the desert of Haroonabad, organic matter is very low so it is not considered good soil. The desert is changing into Death Valley due to high temperature, windstorms and a high rate of evaporation. May and June are the warmest months of this area. There are some plants in this area that come out of the ground after rain and die after producing seeds.

Collection of Ethnobotanical data: The total number of participants in the survey was 85 (Male: 51, Female: 34) and they were interviewed by questionnaire. All participants in the survey were between the ages of 37 and 63 years. The eleven percent (9) of the participants was from the homeopathic field and the rest were from different departments like labors, farming, teachers, trading and house-wife.

Table 1: Demographic data of participants

Sr. No.	Variables	Categories	No. of persons	Percentage
1	Gender	Male	51	60
		Female	34	40
2	Participant profession	Hakim	9	11
		Labor	17	20
		Farming	15	18
		Teachers	13	15
		Trading	19	22
		House-wife	21	25
		3	Educational background	Illiterate
Middle	31	37		
Matric	17	20		
Intermediate	15	18		
Bachelor	8	9		
Specialization	14	16		
4	Age	30-40		4
		40-50	35	41
		50-60	37	43
		60	9	11

Quantitative analysis of Ethnobotanical data

Use Value (UV): The UV was used to explore the importance of any plant species. According to the formula of Phillips *et al.* (1994), the UV was determined.

$$UV = \Sigma U/n$$

U = No. of participants for a given species

n = Total number of participants

Relative Frequency Citation (RFC): According to Vitalini *et al.* (2013), RFC value was calculated

$$RFC = FC / N$$

FC = Number of participants who points out the use of plant

N = Total number of participants

Informant Consensus Factor (ICF): According to Heinrich *et al.* (1998), ICF was calculated

$$ICF = \{(Nur - Nt) / Nur - 1\}$$

Nur = Total number of use informants mentioned for a particular disease category

Nt = Total number of plant species used for a particular disease category

RESULTS

The present study examined the wild flora of Haroonabad, District Bahawalnagar which indicates that the wild flora of this area belongs to 71 species of 28 families. Along with these presented 69 herbs (85%), 5 trees (6%) and 7 shrubs (9%) species by growth habit (Table 4). The most commonly represented family was Poaceae (15 species) followed by Euphorbiaceae (8 species), Asteraceae (7 species) and Amaranthaceae (7 species). *Haloxylon salicornicum* L. and *Cuscuta campestris* Yunk. Were leafless plant species. Annual plants (58%) were more prominent than perennial plants (42%). The studied region supported the Therophytes (59%) and the majority of plants contained the pinnate leaf venation (36%).

The ethnobotanists of Azad Jammu and Kashmir District Kotli collected the 463 plant species belonging to 306 genera and 93 families and reported that the Poaceae family was the dominant among all families [11]. In Pakistan Tolipir National Park, 35 tree species, 19 shrubs, 3 epiphytes, 4 climbers, 75 herbs, 10 ferns, 1 moss and 1 lichen species were recorded. The herbs were dominant in the studied region. The herbs have more importance in the medicinal field due to their high reforestation capacity [5].

Table 3: List of wild flora in Haroonabad, District Bahawalnagar, Pakistan

Sr. No.	Scientific name	Common name	Family	LV	LFS	ST	LS	GH
1	<i>Trianthema portulacastrum</i> L.	Black pigweed	Aizoaceae	PIN	Th	Hr	A	H
2	<i>Zaleya pentendra</i> L.	Biskhapra	Aizoaceae	PIN	Th	Hr	P	H
3	<i>Achyranthes aspera</i> L.	Devil weed	Amaranthaceae	RCL	Th	W	P	H
4	<i>Alternanthera sessilis</i> L.	Gandal booti	Amaranthaceae	PIN	Ch	Hr	P	H
5	<i>Amaranthus viridis</i> L.	jungle cholai	Amaranthaceae	PIN	Th	Hr	A	H
6	<i>Aerva javanica</i> (Burm. f.) Schult.	Bui	Amaranthaceae	PIN	Ch	Hr	P	H
7	<i>Digera muricata</i> L.	Tandla	Amaranthaceae	PIN	Ch	Hr	A	H
8	<i>Haloxylon salicornicum</i> L.	Lana	Amaranthaceae	LL	Ch	W	P	S
9	<i>Suaeda fruticosa</i> (L.) Forsk	Kali lani	Amaranthaceae	RCL	Ha	W	P	S

Contin.....

Sr. No.	Scientific name	Common name	Family	LV	LFS	ST	LS	GH
10	<i>Calotropis procera</i> (Aiton)	Aak	Asclepiadaceae	PIN	Ch	W	P	S
11	<i>Cichorium intybus</i> L.	Kasni	Asteraceae	PIN	Th	Hr	A	H
12	<i>Cirsium arvense</i> L.	Leh	Asteraceae	PIN	Th	Hr	P	H
13	<i>Conyza ambigua</i> L.	Horseweed	Asteraceae	RCL	Th	Hr	A	S
14	<i>Eclipta alba</i> L.	False daisy	Asteraceae	RCL	Ch	Hr	P	H
15	<i>Parthenium</i>	Chatak	Asteraceae	PIN	Th	Hr	A	H

	<i>hysterophorus</i> L.	chandni						
16	<i>Sonchus asper</i> L.	Spiny sow thistle	Asteraceae	PIN	Th	Hr	A	H
17	<i>Sonchus oleraceus</i> L.	Smooth sow thistle	Asteraceae	RCL	Th	Hr	A	H
18	<i>Cordia dichotoma</i> G. Forst	Lasura	Boraginaceae	RCL	Ph	W	P	T
19	<i>Heliotropium indicum</i> L.	Oont chara	Boraginaceae	PIN	Th	Hr	A	H
20	<i>Sisymbrium irio</i> L.	Jangli sarson	Brassicaceae	RCL	Th	Hr	A	H
21	<i>Cleome viscosa</i> L.	Hulhul	Capparidaceae	RCL	Th	Hr	A	H
22	<i>Spergula arvensis</i> L.	Jangli dhania	Caryophyllaceae	RCL	Th	W	A	H
23	<i>Chenopodium ambrosides</i> L.	Worm seed	Chenopodiaceae	RCL	Th	Hr	A	H
24	<i>Chenopodium album</i> L.	Bathu	Chenopodiaceae	PLM	Th	Hr	A	H
25	<i>Chenopodium berlandieri</i> Moq.	Jangli bathu	Chenopodiaceae	RCL	Th	Hr	A	H
26	<i>Chenopodium murale</i> L.	Krund	Chenopodiaceae	RCL	Th	Hr	A	H
27	<i>Convolvulus arvensis</i> L.	Lehli, baily	Convolvulaceae	PIN	Th	Hr	P	H
28	<i>Convolvulus Pluricaulis</i> Choisy.	Makro	Convolvulaceae	PIN	Th	Hr	P	H
29	<i>Citrulus Colocynthis</i> L.	Kor tumma	Cucurbitaceae	PIN	He	W	A	H
30	<i>Cucumis melo</i> L.	Musk-melon	Cucurbitaceae	RCL	T	Hr	A	H
31	<i>Cuscuta campestris</i> Yunk.	Amar bale	Cuscutaceae	LL	Pa	Hr	A	H
32	<i>Cyperus rotundus</i> L.	Chotibhoin, kalooro	Cyperaceae	PAR	Th	Hr	P	H
33	<i>Fimbristylis dichotoma</i> L.	Coco grass	Cyperaceae	PAR	He	Hr	P	H

Contin.....

Sr. No.	Scientific name	Common name	Family	LV	LFS	ST	LS	GH
34	<i>Schoenoplectus mucronatus</i> L.	Rush booti,	Cyperaceae	PAR	Th	Hr	P	H
35	<i>Chrozophora Plicata</i> Vahl.	Giradol	Euphorbiaceae	RCL	Ch	Hr	A	H
36	<i>Euphorbia granulate</i> Orteg.	Hazar dani dodhak	Euphorbiaceae	PAR	Th	Hr	A	H
37	<i>Euphorbia helioscopia</i> L.	Chhatri dodhak	Euphorbiaceae	PIN	Th	Hr	A	H
38	<i>Euphorbia hirta</i> L.	Laldodhak	Euphorbiaceae	RCL	Th	Hr	A	H
39	<i>Euphorbia microphylla</i> Heyne ex. Roth.	Nani dudheli, sandmat	Euphorbiaceae	RCL	Th	Hr	A	H
40	<i>Euphorbia thymifolia</i> L.	Gulf sandmat	Euphorbiaceae	RCL	Th	Hr	A	H

41	<i>Phyllanthus maderaspatensis</i> L.	Kanocha	Euphorbiaceae	PIN	Ch	Hr	A	H
42	<i>Ricinus communis</i> L.	Arind	Euphorbiaceae	PLM	Ph	W	P	S
43	<i>Acacia karoo</i> Hayne.	Pahari keekar	Fabaceae	PIN	Ph	W	P	S
44	<i>Accaia nilotica</i> L.	Keekar	Fabaceae	PIN	Ph	W	P	T
45	<i>Albizia lebbek</i> L.	Sharin	Fabaceae	RCL	Ph	W	P	T
46	<i>Cassia fistula</i> L.	Amaltas	Fabaceae	RCL	Ph	W	P	T
47	<i>Hydrilla verticillata</i> (L.f.) Royle	Jala	Hydrochloritaceae	PAR	Th	Hr	P	H
48	<i>Ocimum Basilicum</i> L.	Niazboo	Limiacae	PIN	Ch	Hr	A	H
49	<i>Oxalis corniculata</i> L.	Khati boti	Oxalidaceae	PIN	Th	Hr	A	H
50	<i>Lathyrus aphaca</i> L.	Jangli matar	Papilionaceae	RCL	Th	Hr	A	H
51	<i>Medicago polymorpha</i> L.	Maina	Papilionaceae	PIN	Th	Hr	A	H
52	<i>Pongamia pinnata</i> L.	Sukhchain	Papilionaceae	PIN	Ph	W	P	T
53	<i>Vicia sativa</i> L.	Revari	Papilionaceae	PIN	Th	Hr	A	H
54	<i>Phyllanthus niruri</i> L.	Gulf leaf flower	Phyllanthaceae	PIN	Ch	Hr	A	H
55	<i>Avena fatua</i> L.	Javi	Poaceae	PAR	Th	Hr	A	H
56	<i>Bromus catharticus</i> Vahl.	Chawli ghass	Poaceae	PAR	He	Hr	P	H
57	<i>Brachiaria ramosa</i> L.	Sudan ghass	Poaceae	PAR	Th	Hr	A	H
58	<i>Cenchrus ciliaris</i> L.	Dhamasa	Poaceae	PAR	He	Hr	P	H
59	<i>Cyanodon dactylon</i> L.	Khabal ghass	Poaceae	PAR	He	Hr	P	H
60	<i>Dactyloctenium aegyptium</i> L.	Madhana ghass	Poaceae	PAR	Th	Hr	A	H

Contin.....

Sr. No.	Scientific name	Common name	Family	LV	LFS	ST	LS	GH
61	<i>Demostachya bipinnata</i> L.	Deep root grass	Poaceae	PAR	He	Hr	P	H
62	<i>Dichanthium annulatum</i> Forssk.	Diaz	Poaceae	PAR	Ch	Hr	P	H
63	<i>Echinochloa colona</i> L.	Jungle rice	Poaceae	PAR	Th	Hr	A	H
64	<i>Echinochloa crusgalli</i> L.	Barnyard grass	Poaceae	PAR	Th	Hr	A	H
65	<i>Imperata cylindrica</i> L.	Nirm dib	Poaceae	PAR	Ge	Hr	P	H
66	<i>Leptochloa chinensis</i> L.	Kallar ghass	Poaceae	PAR	Th	Hr	A	H
67	<i>Polypogon</i>	Dumb ghass	Poaceae	PAR	Th	Hr	A	H

	<i>monspeliensis</i> L.							
68	<i>Sachharum benghalense</i> Retz.	Sarkanda, munj	Poaceae	PAR	He	Hr	P	H
69	<i>Sorghum halepense</i> L.	Baru	Poaceae	PAR	He	Hr	P	H
70	<i>Emex spinosa</i> L.	Trkandi palak	Polygonaceae	RCL	Th	Hr	A	H
71	<i>Rumex dentatus</i> L.	Jangli palak	Polygonaceae	RCL	Ge	Hr	A	H
72	<i>Portulaca oleracea</i> L.	Qulfa, Lonak	Portulacaceae	RCL	Th	Hr	A	H
73	<i>Anagallis arvensis</i> L.	Billi booti	Primulaceae	RCL	Th	Hr	A	H
74	<i>Renunculus muricatus</i> L.	Ghorr summi	Renunculaceae	RCL	Ge	Hr	P	H
75	<i>Datura alba</i> L.	Jimson weed	Solanaceae	RCL	Th	Hr	P	H
76	<i>Physalis minima</i> L.	Rasbari	Solanaceae	RCL	Ch	Hr	A	H
77	<i>Solanum nigrum</i> L.	Mako, Peelak	Solanaceae	RCL	Th	Hr	A	H
78	<i>Withania somnifera</i> L.	Aksin	Solanaceae	PIN	Ch	Hr	P	S
79	<i>Sphenoclea zeylanica</i> Gaertn.	Mirch booti	Sphenocolaceae	PIN	Th	Hr	A	H
80	<i>Corchorus tridens</i> L.	Jangli patsan	Tilaceae	PIN	Th	Hr	A	H
81	<i>Phyla nodiflora</i> L.	Bukkan-booti	Verbenaceae	PIN	Th	Hr	P	H

Legends = LV; Leaf Venation, PIN; Pinnate, RCL; Reticulate, LL; Leafless, PAR; Parallel, PLM; Palmate, LFS; Life Form Spectrum, Th; Therophyte, Ph; Phanerophytes, Ge; Geophytes, Ch; Chaemophytes, Pa; Parasite, He; Hemicryptophytes, ST; Stem type, LS; Life Span, GH; Growth Habitat, Hr; Herbacium, W; Woody, A; Annual, P; Perennial, H; Herb, S; Shurb, T; Tree.

Table 4: Showing the number and percentage of different parameters

Life span					
	No. of Species	Percentage		No. of Species	Percentage
Annual	47	58	Perennial	34	42
Life form spectrum					
	No. of Species	Percentage		No. of Species	Percentage
Phanerophytes	7	8	Chaemophytes	13	17
Geophytes	3	3	Therophytes	48	59
Hemicryptophytes	8	10	Parasite	1	1
Growth habitat					
	No. of Species	Percentage		No. of Species	Percentage
Herbs	69	85	Shrubs	7	9
			Tree	5	6
Leaf venation					
	No. of Species	Percentage		No. of Species	Percentage
Pinnate	29	36	Reticulate	28	34
Parallel	20	25	Palmate	2	2.5
Stem type					
	No. of Species	Percentage		No. of Species	Percentage
Herbaceous	68	84	Woody	13	16

Ethnobotanical data

The Ethnobotanical data of 81 plants were collected from 85 participants through a questionnaire. No one knew the use of 11 plants (*Vicia sativa* L., *Medicago polymorpha* L., *Leptochloa Chinese* L., *Emox spinosa* L., *Sorghum halepense* L., *Cascuta campestris* Yunk., *Schoenoplectus mucronatus* L., *Phyllanthus maderaspatensis* L., *Corchorus tridens* L., *Dichanthium annulatum* Forsk. and *Bromus catharticus* Vahl.) out of 81. The range of UV value was 0.09 (*Chrozophora plicata* Vahl.) to 0.78 (*Acacia nilotica* L.) and RCF value range was 0.03 (*Imperata cylindrica* L.) to 0.95 (*Acacia nilotica* L.). The highest UV value means a lot of people were used this plant for treatment of ailments. The highest value of RCF means that a lot of people were declared that this (given) plant was useful in ayurvedic field. Residents of the studied area use various processes like juice, infusion, ash, extract, decoction, paste, tea, powder and poultice etc. to prepare a recipe for the cure of different illnesses. While making of decoction, Plant parts were boiled in water until the volume of water reduced to ¼ of original volume. Crude extract was gained by crushing and squeezing of plant parts. Some people were crushed the plant parts and smell it for the treatment of different diseases. Infusion in the ayurvedic field was considered good because bioactive components in plants were not degraded [4]. The majority of recipes were prepared from fresh plant parts [10]. The informants thought that the flowering period was best for plant collection because a lot of bioactive compounds were activated at this stage. The use of seeds and roots in the medicinal field is responsible for the extinction of plants [6].

Informants were aware of the side effects of wild plants as well as their benefits. *Trianthema portulacastrum* L. leaves were also used in a salad but a high dose of leaves also caused the paralysis. *Amaranthus viridis* L. was not toxic itself but if they grow in nitrogenous soil then it showed the toxic effects on consumers. The high dose of *Calotropis gigantean* (L.) W.T.Aiton slowed down the heartbeat and responsible for vomiting. *Conyza ambigua* L. caused the skin allergy after touching the plant. *Parthenium hysterophorus* L. also known as toxic wild plants with few benefits. It disturbed the respiratory system of consumers. The high dose of seeds of *Cleome viscosa* L. enhanced the stomach problem. The high concentration of oxalic acid *Chenopodium berlandieri* Moq. Produced the kidney stone. The high dose of *Convolvulus Pluricaulis* Choisy. Suddenly reduced the blood pressure. According to the informants, great caution was needed in using *Citrullus Colocynthis* L. as even a small dose of it caused the kidney problem, stomach disease and even death. The seed coat of *Ricinus communis* L. contained the poisonous compounds which caused the stomach ailments and high dose of oil can caused the death. The *Acacia nilotica* L. also needed a lot of caution in using it. It also had some side effects. If it used indiscriminately, it can be caused liver and stomach diseases. The un-ripened fruit of *Solanum nigrum* L. also showed the toxic effect.

According to researchers, the plants of Euphorbiaceae, Fabaceae, Asteraceae and Apocynaceae were considered toxic because majority of plants was contained a saponins, glycosides, steroids, alcohol, cyanogenic glycosides, resins and selenium etc. [3]. Herbs or wild plants were not beneficial or toxic. The use of plants made them beneficial or toxic. The dose of medicines varied from child to old. Dose was taken thrice or twice a day depending upon the

nature or situation of patients. The people used the inappropriate techniques for the preparation of medicines which showed the toxic or side effects on the health of consumers [12].

Table 5: Ethnobotanical data of wild flora of selected area

Species name	Part used	Recipes with ailments	UV	RFC
<i>Trianthema portulacastrum</i> L.	Leaves,	Leaves	0.35	0.13
	Roots,	*Powder and fresh leaves (odema, dropsy and jaundice)		
	Whole plant	Roots *Powder (liver, asthma and veneral discharge)		
		Whole plant *Powder (alcoholic poisoning, veneral discharge, heart disease and piles)		
<i>Zaleya pentendra</i> L.	Leaves,	Leaves	0.23	0.09
	Whole plant	*Juice (stomach problem)		
		Whole plant *Powder (urinary infection)		
<i>Achyranthes aspera</i> L.	Seeds,	Seeds	0.31	0.07
	Flowers,	*Snake bite		
	Roots,	Flowers		
	Whole plant	*Rubbed with sugar and made medicine (dog bite)		
		Roots *Powder (stomach, cholera)		
		Leaves *Paste (scorpion bite, insect sting paralysis and nervous system disorder)		
		Whole plant *Powder (washing soda, malaria and cleaning teeth)		
		*Extract (diarrhea, dysentery and skin diseases)		
		Roots		
		*Juice (fever, cold, cough and blood dysentery)		
<i>Alternanthera sessilis</i> L.	Roots,	Roots	0.51	0.23
	Whole plant	*Juice (fever, cold, cough and blood dysentery)		
		Whole plant *Paste (wounds, draw out spines from body and cooling agent)		
		*Baked with corn flour (menstrual disorder and stomach disease)		
		*Dry plant mixed with salt (Stop excess bleeding and blood vomiting)		

<i>Spergula arvensis</i> L.	Seeds	*Making bread	0.14	0.16
<i>Chenopodium ambrosoides</i> L.	Whole plant	*Tea (inflammatory and lung infection)	0.16	0.09

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Amaranthus viridis</i> L.	Leaves, Roots, Whole plant	Leaves *Juice (heart disease and eye wash to prevent infections) *Powder (inflammation) Roots *Juice (inflammation of urinary bladder, constipation and dysentery) Whole plant *Powder (soap making)	0.33	0.28
<i>Aerva javanica</i> (Burm. f.) Schult.	Seeds, Flowers, Roots, Leaves	Seeds *Boiled seeds (mouth disease) Flowers *Dry and fresh are used to stop the bleeding and repair the damaged cell. Roots *Juice (eye washing) Leaves *Paste (inflammation)		
<i>Digera Arvensis</i> L.	Stems, Seeds, Flowers	Stems *Chewing of stem improve the digestive system Seeds *Urinary disorder Flowers *Dry (Urinary disease)	0.15	0.19
<i>Haloxylon salicornicum</i>	Leaves	Leaves *Tea (minimize the pregnancy pain) *Fresh juice (eye infection) Whole plant *Ash of whole plant (wounds)	0.17	0.21
<i>Suaeda fruticosa</i> (L.) Forsk	Leaves	Leaves *Paste (ringworm infection and skin allergy)	0.19	0.16
<i>Calotropis gigantean</i> (L.)	Barks, Root, Leaves	Barks *Smoke is inhaled (respiratory disease and	0.42	0.56

W.T.Aiton	asthma) *Dry powder mix in water (stomach and heart problem) Roots *Ash of dry roots (skin allergy) Leaves *Leaf juice rubbed on scorpion sting
-----------	--

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Cichorium intybus</i> L.	Leaves, Flowers, Whole plant	Leaves and flower *Paste (wounds) Whole plant *Juice (diarrhea and stomach problem)	0.34	0.26
<i>Cirsium arvense</i> L.	Stems, Roots	Stems *Chewing (toothache) Roots *Ash (wounds)	0.16	0.16
<i>Conyza ambigua</i> L.	Roots, Whole plant	Roots *Root tea (menstrual disorder) Whole plant *Tea (alcoholic poisoning) *Paste (piles) *Steam (enhance sneezing during cold)	0.18	0.14
<i>Eclipta alba</i> L.	Leaves	*Chewing few leaves daily (eye health) *Extract (high blood pressure)	0.13	0.24
<i>Parthenium hysterophorus</i> L.	Whole plant	*Decoction (diarrhea, urinary infection, fever and malaria)	0.20	0.31
<i>Sonchus asper</i> L.	Whole plant	*Ash (wounds)	0.11	0.17
<i>Sonchus oleraceus</i> L.	Leaves	*Juice (eye disease) *Paste (inflammation)	0.23	0.13
<i>Cordia dichotoma</i> G. Forst	Seeds, Barks, Fruits, Leaves	Seeds *Powder apply on skin allergy Barks *Paste (swelling) *Maswak (teeth disease and headache) Fruits *Paste (skin allergy) Leaves *Juice (cooling)	0.35	0.47

<i>Heliotropium indicum</i> L.	Leaves, Whole plant	Leaves *Infusion (asthma) *Paste (insect sting) Whole plant *Decoction (thrush and diabetes) *Boiled and rubbed on heat rash	0.37	0.18
<i>Chenopodium berlandieri</i> Moq.	Whole plant	*Cooked (gout, kidney stone and arthritis)	0.30	0.47
<i>Chenopodium murale</i> L.	Seeds	*Powder (making bread with wheat and kidney stone)	0.41	0.68

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Sisymbrium irio</i> L.	Leaves, Seeds, Whole plant	Leaves *Infusion (throat and chest infection) Seeds *Seed in water (asthma) Whole plant *Paste (inflammation)	0.59	0.19
<i>Cleome viscosa</i> L.	Leaves, Whole plant	Leaves *Paste (wounds) Whole plant *Juice (stimulate the appetite) *Decoction (improve the digestive system and enhance hunger)	0.22	0.11
<i>Chenopodium album</i> L.	Leaves, Seeds, Stems, Roots, Whole plant	Leaves *Paste (swelling) Seeds *Chewing (urinary infection) Stems *Juice (freckles) Roots *Juice (blood dysentery) Whole plant *Decoction (sunstroke and teeth disease)	0.68	0.87
<i>Chenopodium berlandieri</i> Moq.	Whole plant	*Cooked (gout, kidney stone and arthritis)	0.30	0.47
<i>Chenopodium murale</i> L.	Seeds	*Powder (making bread with wheat and kidney stone)	0.41	0.68
<i>Convolvulus arvensis</i> L.	Flower, Leaves,	Flower *Tea (fever)	0.23	0.12

	Whole plant	Leaves *Tea (reduce the excessive menstrual)		
<i>Convolvulus Pluricaulis</i> Choisy.	Whole plant	Whole plant *Juice of vine with water (Liver infection) *Decoction with cumin and milk (enhance memory) *Juice (headache) *Powder (high blood pressure)	0.45	0.34
<i>Citrullus Colocynthis</i> L.	Whole plant	*Juice diluted with water (skin infection)	0.47	0.39
<i>Chrozophora Plicata</i> Vahl.	Whole plant	*Paste (wound) *Juice (purification of blood)	0.09	0.03

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Cucumis melo</i> L.	Fruits, Seeds, Leaves	Fruits *Fruit (cooling) Seeds *Ground and take with water (improve digestive system) Leaves *Fresh or dry leaves are used to release hernias	0.19	0.25
<i>Cyperus rotundus</i> L.	Roots, Tubers	Roots *Dry Roots (digestive and menstrual problem) *Dry Root mixed with black pepper (stomach problem) Tubers *Dry tuber powder (prevent teeth decay)	0.14	0.12
<i>Fimbristylis dichotoma</i> L.	Roots, Leaves	Roots *Crushed roots (aphrodisiac) Leaves *Leaves as a poultice (fever)	0.16	0.08
<i>Euphorbia granulate</i> Orteg.	Latex	*Latex purify the blood and externally used on sting bite or scorpion	0.31	0.25
<i>Euphorbia helioscopia</i> L.	Leaves	*Extract used for HIV-1 and hepatitis C	0.23	0.18
<i>Euphorbia hirta</i> L.	Leaves, Stems, Whole plant	Leaves *Infusion (asthma) Stems *Infusion (asthma) Whole plant	0.45	0.19

<i>Euphorbia microphylla</i> Heyne ex. Roth.	Whole plant	*Paste (sting bite, skin infection, inflammation and fungal infection) *Decoction (foot athletes and fungal infection) *Infusion (asthma) *Decoction (improve digestive system)	0.17	0.21
<i>Euphorbia thymifolia</i> L.	Leaves, Whole plant	Leaves *Decoction (kidney disease) *Paste (headache) Whole plant *Decoction (diarrhea, lung problem, eye wash and venereal diseases)	0.19	0.08
<i>Hydrilla verticillata</i> (L.f.) Royle	Whole plant	*Chewing or juice (Nervous system disorder, build blood cells and regeneration of skin)	0.23	0.09

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Ricinus communis</i> L.	Fruits	Oil *Castor oil helpful for reducing constipation when taken by mouth *Dry eyes cure by castor oil *Castor oil removed the dandruff in hair *Few drops of castor oil used for better vision	0.54	0.89
<i>Acacia karoo</i> Hayne.	Stems, Whole plant	Stems *Extract (loose motion, diarrhea and urinary bladder pain) *Chewing (oral ailment) Whole plant *Whole plant juice used for wound washing	0.66	0.93
<i>Oxalis corniculata</i> L.	Leaves, Whole plant	Whole plant *Juice (stomach, influenza, urinary disease, insect bite and scurvy) Leaves *Juice (insect sting and skin cramps)		
<i>Accaia nilotica</i> L.	Barks, Flowers, Leaves, Seeds, Whole plant	Barks *Chewing (reduce the teeth lose and stop the bleeding) *Gums (skin irritation, inflammation and Diabetes) *Powder (toothaches) *Boiled with water and wash the wounds Flowers *Fresh (loose motion)	0.78	0.95

		Leaves *Juice (eye washing) *Leave (wounds) Seeds *2 gm seeds with warm water (high blood pressure) Whole plant *Decoction (diarrhea)		
<i>Albizia lebbek</i> L.	Barks, Flowers, Seeds, Leaves	Barks *Chewing (diarrhea and piles) Flowers *Paste (skin infection) Seeds *Seeds with mishri (weakness) Leaves *Grind with honey and water (urinary problem)	0.70	0.98
Contin.....				
Species name	Part used	Recipes with ailments	UV	RFC
<i>Cassia fistula</i> L.	Leaves, Pods, Barks	Leaves *Extract (skin pathogen) Pods *Decoction (malaria, diabetes and kidney stone) Bark *Paste (skin disease) *Decoction (washing wounds)	0.61	0.84
<i>Ocimum Basilicum</i> L.	Leaves, Seeds, Whole plant	Leaves *Juice (Influenza and enhance digestion) *Paste (skin allergy and snake sting) Seeds *Infusion (diarrhea) Whole plant *Juice (digestive and nervous system disorder)	0.45	0.59
<i>Lathyrus aphaca</i> L.	Seeds	*Chewing (teeth diseases)	0.19	0.09
<i>Pongamia pinnata</i> L.	Seeds, Leaves, Stems, Roots	Seed *Oil stomach and liver disease *Paste joint disorder and sores Leaves *Decoction (cough) *Paste (skin infection and stop bleeding)	0.44	0.81

		Stems		
		*Maswak (Stop spleen enlargement)		
		Roots		
		*Juice (ulcer and toothaches)		
<i>Phyllanthus niruri</i> L.	Roots, Leaves	Roots	0.18	0.08
		*Juice (bladder stone)		
		*Infusion (chronic dysentery and cold)		
		Leaves		
		*Infusion (dysentery, stomach problem and cold)		
		*Paste mixed with salt (wounds)		
<i>Avena fatua</i> L.	Seeds	*Seeds are used as a cereal for nourishment	0.30	0.45
<i>Brachiaria ramosa</i> L.	Roots, Whole plant	Roots	0.23	0.18
		*Paste (kidney pain)		
		Whole plant		
		*Ash(insect sting)		
<i>Cenchrus ciliaris</i> L.	Whole plant	*Juice (kidney problem and tumor)	0.15	0.12
		*Paste (wound)		

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Cyanodon dactylon</i> L.	Leaves, Whole plant	Leaves *Paste (wound)	0.37	0.15
		Whole plant *Decoction (cancer, cough, headache, dropsy, stone, snake sting, wound and toothaches)		
<i>Dactyloctenium aegyptium</i> L.	Leaves	*Infusion (accelerate childbirth)	0.29	0.17
<i>Demostachya bipinnata</i> L.	Whole plant	*Decoction (dysentery)	0.12	0.14
<i>Echinochloa colona</i> L.	Stems	*Decoction (dysentery)		
<i>Echinochloa crusgalli</i> L.	Roots	*Paste (wounds)	0.16	0.13
<i>Imperata cylindrica</i> L.	Flowers, Roots, Whole plant	Flowers *Paste (wound)s *Decoction (urinary infection and fever)	0.10	0.03
		Roots *Decoction (digestive system diseases)		
		Whole plant *Extract (cancer)		
	Whole plant	*Infusion (heart problem)	0.23	0.05

<i>Polypogon monspeliensis</i> L.				
<i>Sachharum benghalense</i> Retz.	Leaves, Stems	Leaves *Juice (eye wash) *Decoction (urinary infection) Stems *Juice (sore throat)	0.15	0.17
<i>Rumex dentatus</i> L.	Roots, Leaves, Whole plant	Roots *Decoction (stomach problem and intestinal parasite) *Paste (swelling) Leaves *Juice (headache) *Paste (wounds) Whole plant *Decoction (reduce body pain)	0.22	0.23
<i>Portulaca oleracea</i> L.	Leaves Whole plant	Leaves *Juice (heart disease, cough and earaches) *Tea (stomach disease and headache) Whole plant *Extract (muscle relaxation and wounds) *Juice (skin diseases and insect bite)	0.34	0.25

Contin.....

Species name	Part used	Recipes with ailments	UV	RFC
<i>Anagallis arvensis</i> L.	Whole plant	*Infusion (skin diseases and liver diseases) *Paste (skin itches and warts)	0.16	0.16
<i>Renunculus muricatus</i> L.	Fruits	*Fresh or juice (asthma and fever)	0.28	0.26
<i>Datura alba</i> L.	Leaves, Seeds	Leaves *Paste of roasted leaves (reduce full body pain) *Smoke (asthma) *Juice (ear disease) Seeds *Oil (stimulate hair growth)	0.10	0.04
<i>Physalis minima</i> L.	Fruits, Leaves, Roots, whole plant	Fruits *Fresh (enhance appetite) Leaves *Paste (headache and skin rash) Roots *Extract (fever) *Decoction (diabetes)	0.21	0.14

		Whole plant		
		*Extract (cancer)		
<i>Withania somnifera</i> L.	Whole plant	*Juice (tiredness. High sugar level in blood and high cholesterol level)		
<i>Solanum nigrum</i> L.	Fruits,	Fruits	0.47	0.57
	Whole plant	*Juice (teeth diseases)		
		*Fresh (eye disease and fever)		
		*Paste (headaches and skin allergy)		
		Whole plant		
		*Paste (wounds)		
<i>Sphenoclea zeylanica</i> Gaertn.	Leaves	*Paste (sting of poisonous animals)	0.17	0.19
<i>Phyla nodiflora</i> L.	Leaves,	Leaves	0.22	0.33
	Whole plant	*3-4 fresh leaves (piles)		
		*Paste (infected skin by ulcer)		
		Whole plant		
		*Whole plant juice (cough and fever)		

Part used: The Ethnobotanical data related to part used of 70 species was displayed because 11 plant species had no Ethnobotanical data and 57% whole plant was used in the ayurvedic field followed by leaves (53%) and roots (26%). According to the data cited by the informants, the %age of using pods (*Cassia fistula* L.), tubers (*Cyperus rotundus* L.) and latex (*Euphorbia granulate* Orteg.) was minimal because they were used only by one plant. The use of leaves other than other plant plants in the medicinal field was good because leaves are easily collected and sustainable method in ecology with no fear of extinction [7].

Table 6: Percentage of plant part used

Part used	No. of Species	Percentage	Part used	No. of Species	Percentage
Leaves	37	53	Roots	18	26
Seeds	16	23	Fruits	6	8
Flowers	8	11	Stems	8	11
Barks	5	7	Whole plant	40	57

Informant consensus factor (ICF): ICF was used to focus on the consistency of data related to particular disease categories. The disease was divided into 15 categories based on the recorded ailments by participants. The range of ICF was recorded from 0.45 to 0.78. The lowest ICF value was observed in hormonal disorder (0.45) and the highest value was shown by respiratory disease (0.78). The highest ICF value predicts that the high %age of participants was used plants

for an illness of a comprehensive category. The local people of Haroonabad were used *Euphorbia helioscopia* L. leaves to cure HIV and hepatitis C and *Withiana somnifera* L. was used to treat the high sugar and cholesterol level. The leaves of *Oxalis corniculata* L. were used to treat the scurvy ailment and *Cynodon dactylon* L. was used in cancer treatment.

Table 7: ICF of recorded plant species with respect to different ailments.

Sr. No.	Disease category	No. of use reports	Percentage of use reports	No. of species	Percentage of use species	*ICF
1	Respiratory diseases	41	48	10	12	0.78
2	Nervous system disorder	7	8	4	5	0.50
3	Skin infection	61	72	24	30	0.61
4	Urinary diseases	19	22	10	12	0.50
5	Hormonal disorder	10	12	6	7	0.45
6	Heart diseases	11	13	5	6	0.60
7	Digestive system diseases	17	20	9	11	0.50
8	Stomach diseases	43	51	11	14	0.76
9	Liver diseases	15	18	3	3	0.85
10	Wounds	70	82	18	22	0.75
11	Insect sting	31	36	11	14	0.67
12	Muscle disorder	13	15	3	3	0.83
13	Stone	27	32	7	9	0.77
14	Diabetes	9	11	4	5	0.63
15	Eye, ear, throat and mouth disease	29	34	13	16	0.57

CONCLUSION

The biodiversity of any area indicates the importance of the area. Poaceae and Therophyte species were more prominent in the study area. Wild flora maintains the ecosystem and sustainability in the environment and they also had a major role in the medical field. Present research discovered the medicinal plants that mostly used by local inhabitants to cure various ailments. It tries to attract the attention towards the conservation policy of wild plants. The wild plants in Haroonabad have very importance in the homeopathic medical field but biodiversity was not high.

RECOMMENDATIONS

1. The native people of studied region have used traditional botanical knowledge to heal the various ailments but the accessibility of plants is atypical. Consequently, the traditional healers and the native people should be awake on how to use plants for a variety of objectives and the traditional healers should cultivate some plants in their home gardens.
2. There are some medicinal plants used to treat different diseases. But the chemical components of these plants are not well known. Therefore, research on chemical analysis of these plants should be performed.
3. There is a need to authorize the indigenous communities and make sure their active participation in sustainable harvesting and conservation of natural resources. Different

universities should collaborate with indigenous communities and recognize them as 'knowledge site' on a particular subject to uphold their status and conserve its knowledge.

REFERENCES

1. Anonymous, 2009. Flora of Ziarat: Ethnobotanic and Medicinal Importance, pp. 71.
2. Anyinam C. (1995). "Ecology and Ethnomedicine: Exploring Links between Current Environment Crisis and Indigenous Medical Practices," *Soc. Sci. Med.*, 40(3): 321-329.
3. Barla A, Bırman H, Kültür Ş and Öksüz S. (2006). Secondary metabolites from *Euphorbia helioscopia* and their vasodepressor activity. *Turk. J. Chem.* 30(3): 325-332.
4. Dike IP, Obembe OO and Adebisi FE. (2012). Ethnobotanical survey for potential anti-malarial plants in south-western Nigeria. *J. Ethnopharmacol.*, 144(3): 618-626.
5. Faiz A, Ghufarn MA, Mian A and Akhtar T. (2014). Floral diversity of Tolipir National Park (TNP), Azad Jammu and Kashmir. *Biologia*, 60(1): 43-55.
6. Ghimire SK, Gimenez O, Pradel R, McKey D and Aumeeruddy-Thomas Y. (2008). Demographic variation and population viability in a threatened Himalayan medicinal and aromatic herb *Nardostachys grandiflora*: matrix modelling of harvesting effects in two contrasting habitats. *J. Appl. Ecol.*, 45(1): 41-51.
7. Giday M, Asfaw Z and Woldu Z. (2010). Ethnomedicinal study of plants used by Sheko ethnic group of Ethiopia. *J. Ethnopharmacol.*, 132 (1): 75-85.
8. Heinrich M, Ankli A, Frei B, Weimann C and Sticher O. (1998). Medicinal plants in Mexico: Healers' consensus and cultural importance. *Soc. Sci. Med.*, 47(11): 1859-1871.
9. Ilyas M, Qureshi R, Shaheen H, Ahmad W and Munir M. (2014). Phytodiversity and plant life of Khanpur Dam, Khyber Pakhtunkhwa, Pakistan. *Pak. J. Bot.*, 46(3): 841-849.
10. Jima TT and Megersa M. (2018). Ethnobotanical study of medicinal plants used to treat human diseases in berbere district, bale zone of oromia regional state, south east Ethiopia. *Evid.-Based Complementary Altern. Med.*, pp 1-16.
11. Khan AM, Qureshi R, Qaseem MF, Munir M, Ilyas M and Saqib Z. (2015). Floristic checklist of district kotli, azad jammu and kashmir. *Pak. J. Bot.*, 47(5): 1957-1968.
12. Lee S, Xiao C and Pei S. (2008). Ethnobotanical survey of medicinal plants at periodic markets of Honghe Prefecture in Yunnan Province, SW China. *J. Ethnopharmacol.*, 117(2): 362-377.
13. Noman A. (2003). Influence of different doses of nitrogen fertilizer on ajwain. M.Sc. Dissertation. pp-1. Univ. of Agric. Faisalabad. Pakistan.
14. Phillips O, Gentry AH, Reynel C, Wilkin P, Galvez-Durand B. (1994). Quantitative ethnobotany and Amazonian conservation. *Conserv. Biol.*, 8(1): 225-248.
15. Shad AA, Shah HU and Bakht J. (2013). Ethnobotanical assessment and nutritive potential of wild food plants. *J. Anim. Plant Sci.*, 23(1): 92-99.
16. Shah A, Poudel RC, Ishtiaq M, Sarvat R, Shahzad H, Abbas A, Shoaib S, Nuzhat R, Noor UD, Mahmooda H, Summaya A, Ifra and Ihsan U. (2019). Ethnobotanical study of medicinal plants of namal valley, salt range, Pakistan. *Appl. Ecol. Environ. Res.*, 17(2):4725-4805.

17. Shinwari ZK and Qaiser M. (2011). Efforts on conservation and sustainable use of medicinal plants of Pakistan. *Pak. J. Bot.*, 43(1): 5-10.
18. Vitalini S, Iriti M, Puricelli C, Ciuchi D, Segale A and Fico G. (2013). Traditional knowledge on medicinal and food plants used in Val San Giacomo (Sondrio, Italy) -An alpine ethnobotanical study. *J. Ethnopharmacol.*, 145(2): 517-529.